



Additions to Philippine Slender Skinks of the *Brachymeles bonitae* Complex (Reptilia: Squamata: Scincidae) II: a new species from the northern Philippines

CAMERON D. SILER^{1,2}, DREW R. DAVIS³, ELYSE S. FREITAS¹, NICHOLAS A. HURON¹,
AARON D. GEHEBER¹, JESSA L. WATTERS², MICHELLE L. PENROD¹, MONICA PAPEŞ⁴,
ANDREW AMREIN¹, ALYSSA ANWAR¹, DONTAE COOPER¹, TUCKER HEIN¹,
ANNALISA MANNING¹, NEERAL PATEL¹, LAUREN PINAROC¹,
ARVIN C. DIESMOS⁵, MAE L. DIESMOS⁶, CARL H. OLIVEROS⁷ & RAFE M. BROWN⁸

¹Department of Biology, University of Oklahoma, 730 Van Vleet Oval, Norman, OK 73019, USA. E-mails: camsiler@ou.edu, aaron.d.geheber-1@ou.edu, michelle.l.penrod-1@ou.edu, elysefreitas@gmail.com, nahuron@ou.edu, a.w.amrein@gmail.com, alyssa.anwar@ou.edu, dontae.cooper@ou.edu, annalisa.r.manning-1@ou.edu, neeral.s.patel-1@ou.edu, lpinaroc2@yahoo.com

²Sam Noble Oklahoma Museum of Natural History, University of Oklahoma, 2401 Chautauqua Avenue, Norman, OK 73072, USA. E-mails: jwatters@ou.edu

³Department of Biology, University of South Dakota, 414 East Clark Street, Vermillion, SD 57069, USA. E-mail: drew.davis@usd.edu

⁴Department of Integrative Biology, Oklahoma State University, 501 Life Sciences West, Stillwater, OK, 74074 USA. E-mail: papes@okstate.edu

⁵Herpetology Section, Zoology Division, Philippine National Museum, Rizal Park, Burgos Street, Manila, Philippines. E-mail: arvin.diesmos@gmail.com

⁶University of Santo Tomas, Espana Boulevard, Manila, Philippines. E-mail: maediesmos@gmail.com

⁷Department of Biological Sciences, Louisiana State University, 220 Life Sciences Building, Baton Rouge, LA 70803, USA. E-mail: carloliveros@gmail.com

⁸Biodiversity Institute and Department of Ecology and Evolutionary Biology, University of Kansas, 1345 Jayhawk Boulevard, Lawrence, KS 66045, USA. E-mail: rafe@ku.edu

Abstract

We describe a new digitless scincid lizard of the genus *Brachymeles* from northern Luzon and Camiguin Norte islands in the Philippines. This species belongs to the *Brachymeles bonitae* Complex, and both molecular and morphological data confirm that this species is distinct from all other congeners. Formerly considered to be a single widespread species, this group of species has been the focus of recent systematic reviews. Here we describe a new species in the *B. bonitae* Complex, recognized currently to constitute five species. *Brachymeles ilocandia* **sp. nov.** is the second digitless and the seventeenth non-pentadactyl species in genus. The description of this species brings the total number of species in the genus to 40, and provides new insight into unique distribution patterns of species of the northern Philippines.

Key words: Babuyan Island Group, biodiversity, ecological niche modeling, endemism, faunal region, fossorial, Luzon Island, non-pentadactyl, pentadactyl

Introduction

The genus *Brachymeles* Duméril & Bibron includes a moderate radiation of semi-fossorial (burrowing) lizards in the family Scincidae that is distributed throughout the Philippine archipelago and in limited areas of mainland and insular Southeast Asia. This genus has been the focus of numerous recent taxonomic studies, and as a result of several phylogenetic studies and examination of museum collections, alpha diversity within *Brachymeles* has increased dramatically over the last five years, from 15 to 39 species (for review, see Davis *et al.* 2014; Geheber *et al.* 2016). The majority of species of *Brachymeles* are found exclusively in the Philippines; the two exceptions include *B. miriamae* Heyer from Thailand and *B. apus* Hikida from Borneo (Heyer 1972; Hikida 1982). Though the morphology and evolutionary history of *Brachymeles* are both well represented in the literature, the ecology of

these species is not well known (Davis *et al.* 2014). Species prefer dry habitats and have been found in leaf litter, loose soil, or under decaying logs (Siler *et al.* 2009, 2010a,b, 2011a,b,c,d, 2012a; Davis *et al.* 2014). Most species are found in disturbed secondary growth lowland habitats; however, several species (*B. apus*, *B. elerae* Taylor, *B. muntingkamay* Siler, Rico, Duya & Brown, and *B. wrighti* Taylor) inhabit mid- to high-elevation habitats (Siler 2010; Siler *et al.* 2009, 2011d).

With the exception of differences in limb and digit morphology, species in the genus *Brachymeles* have conserved body plans and pigmentation patterns. All species are slender, elongate lizards possessing relatively homogeneous brown scale coloration. Interestingly, the clade represents one of only a handful of scincid genera to possess a full spectrum of digit and limb states, from species with more robust, pentadactyl limbs to those with externally limbless bodies (Siler & Brown 2010, 2011; Siler *et al.* 2011a, 2012a; Davis *et al.* 2014). Of the 39 currently recognized species of *Brachymeles*, 18 are pentadactyl (*B. bicolor* Gray, *B. bohollensis* Brown & Rabor, *B. boulengeri* Taylor, *B. gracilis* Fischer, *B. hilong* Brown & Rabor, *B. kadwa* Siler & Brown, *B. makusog* Siler, Diesmos & Brown, *B. mindorensis* Brown & Rabor, *B. orientalis* Brown & Rabor, *B. samad* Siler, Jones, Diesmos, Diesmos & Brown, *B. schadenbergi* Fischer, *B. suluensis* Taylor, *B. talinis* Brown, *B. taylori* Brown, *B. tiboliorum* Siler, Jones, Diesmos, Diesmos & Brown, *B. tungaoi* Siler & Brown, *B. vindumi* Siler & Brown, and *B. vulcani* Siler, Jones, Diesmos, Diesmos & Brown), 16 are non-pentadactyl and have limbs that are reduced in size and digit number (*B. bicolandia* Siler, Fuiten, Jones, Alcalá & Brown, *B. bonitae* Duméril & Bibron, *B. brevidactylus* Siler, Fuiten, Jones, Alcalá & Brown, *B. cebuensis* Brown & Rabor, *B. cobos* Siler, Fuiten, Jones, Alcalá & Brown, *B. elerae*, *B. isangdaliri* Davis, Feller, Brown & Siler, *B. libayani* Siler, Fuiten, Jones, Alcalá & Brown, *B. mapalanggaon* Davis, Feller, Brown & Siler, *B. muntingkamay*, *B. paeorum* Siler, Fuiten, Jones, Alcalá & Brown, *B. pathfinderi* Taylor, *B. samarensis* Brown, *B. ligtas* Geheber, Davis, Watters, Penrod, Feller, Davey, Ellsworth, Flanagan, Heitz, Moore, Nguyen, Roberts, Sutton, Sanguila, Brown & Siler, *B. tridactylus* Brown, and *B. wrighti*), and five are completely limbless (*B. apus*, *B. lukbani* Siler, Balete, Diesmos & Brown, *B. minimus* Brown & Alcalá, *B. miriamae*, and *B. vermis* Taylor) (Davis *et al.* 2014; Geheber *et al.* 2016). Non-pentadactyl, reduced-limbed species represent a sizeable portion of the morphological diversity observed within the genus. Of these species, one is digitless (*B. mapalanggaon*), one is monodactyl (*B. isangdaliri*), four are bidactyl (*B. bicolandia*, *B. brevidactylus*, *B. cobos*, and *B. samarensis*), four are tridactyl (*B. libayani*, *B. muntingkamay*, *B. paeorum*, and *B. tridactylus*), and two are tetradactyl (*B. elerae*, and *B. wrighti*) (Davis *et al.* 2014). Additionally, three species have unequal numbers of fingers and toes, with *B. cebuensis* having three fingers and two toes, *B. ligtas* having two fingers and zero toes, and *B. pathfinderi* with five fingers and four toes (Siler *et al.* 2011c). Finally, *Brachymeles bonitae* sensu stricto represents an instance of both cases: some individuals have digitless limbs while others have two fingers and one toe (Davis *et al.* 2014).

Philippine species of *Brachymeles* form a monophyletic clade, suggesting *in situ* diversification of the genus across the archipelago (Siler & Brown 2011; Siler *et al.* 2011a). Hypotheses for the high levels of Philippine diversity of plants and animals have focused on the influence of changing sea levels on species diversification (Brown *et al.* 2013). During Pleistocene glacial cycles, decreases in the sea level led to the formation of Philippine Aggregate Island Complexes (PAICs; Brown & Guttman 2002; Brown & Diesmos 2002, 2009; Fig. 1A) in which adjacent islands separated by shallow seas were connected by land bridges, allowing for faunal exchange and gene flow between islands within a single PAIC (Brown & Diesmos 2002; Brown *et al.* 2002, 2009, 2013). Although this process likely contributed to the development of distinct faunal regions within the Philippines, studies suggest that species diversification patterns in *Brachymeles* do not follow predicted PAIC-based diversification patterns, with evidence suggesting a number of overseas dispersal events may have also taken place during the radiation of this group throughout the archipelago (Siler *et al.* 2011a). Furthermore, studies on radiations of other Philippine reptiles (Linkem *et al.* 2010, 2011; Siler *et al.* 2010c, 2012b, 2014; Welton *et al.* 2013, 2014; Brown & Siler 2013) have also partially or fully rejected PAIC formation and fragmentation events in the generation and maintenance of species diversity. These studies suggest that additional diversification mechanisms need to be explored to accurately assess and understand the processes that have resulted in the country's diversity of *Brachymeles*. Regardless of the mechanisms involved, amphibian and reptile diversity in the archipelago is largely endemic (Brown *et al.* 2013; Diesmos *et al.* 2015).

The genus *Brachymeles* was first described in 1839 by Duméril and Bibron for the species *B. bonitae*, a small, elongate, limb-reduced species from Manila, on Luzon Island. To date, 11 published reviews or descriptions of the genus or species groups within the genus have improved our understanding of this unique radiation of burrowing

lizards (Taylor 1917; Brown 1956; Brown & Rabor 1967; Brown & Alcala 1980; Siler & Brown 2010; Siler *et al.* 2011a,b,c,d, 2012a; Davis *et al.* 2014). Furthermore, much of the recent taxonomic resolution has been fueled by continued field surveys across the Philippines and the incorporation of genetic data into studies exploring species-level diversity in the genus. Guided by robust phylogenetic estimates, subsequent assessments of morphological diversity revealed many historically recognized widespread species were actually complexes of numerous distinct evolutionary lineages (Siler & Brown 2010, 2011; Siler *et al.* 2011a, 2012a; Davis *et al.* 2014; Geheber *et al.* 2016). As a result, most of these complexes have now been appropriately revised to formally recognize all unique evolutionary lineages (four species of the *B. boulengeri* Complex [Siler & Brown 2010], five species of the *B. bonitae* Complex [Davis *et al.* 2014; Geheber *et al.* 2016], seven species of the *B. gracilis* Complex [Siler *et al.* 2012a], nine species of the *B. samarensis* Complex [Siler *et al.* 2011b], two species of the *B. schadenbergi* Complex [Siler & Brown 2010], and three species of the *B. talinis* Complex [Siler & Brown 2010]).




FIGURE 1. (A) Map of the Philippines islands, with labels for islands discussed in the study, the five recognized major PAICs, major island groups, and additional deep-water islands provided for reference. Current islands in the Philippines are shown in medium grey to black; light grey areas enclosed in black 120 m bathymetric contours indicate the hypothesized maximum extent of land during the mid- to late Pleistocene. Vouchered localities for *Brachymeles ilocandia* sp. nov. marked by red circles. (B) Mapped output of Maxent results as the mean ecological niche model of suitable climatic conditions for *Brachymeles ilocandia* sp. nov. projected across the known distribution of the genus throughout the Philippines. Values corresponding to the likelihood of climatic suitability are included for scale and interpretation of heat map color-coding. Values are logistic-transformed for ease of interpretation on a scale of 0–1, according to the default Maxent output settings (ENM analysis and projection performed by NAH and MP).

With the exception of the historically recognized distribution of *B. talinis* spanning from the southernmost islands of the Mindanao Faunal Region in the Philippines to the Babuyan Island Complex north of the Luzon Faunal Region (Fig. 1A; Siler & Brown 2010; Siler *et al.* 2011a), *B. bonitae* was recognized as having one of the largest distributions of any species within the genus, with populations found throughout much of the central and northern Philippines and spanning multiple PAICs (Siler *et al.* 2011a; Davis *et al.* 2014; Geheber *et al.* 2016). However, recent investigations focused on this widespread species have clarified the distributions of true *B. bonitae* and the closely related tridactyl species *B. tridactylus*, and recognized three additional lineages within the species complex (Davis *et al.* 2014; Geheber *et al.* 2016). Currently, the *B. bonitae* Complex comprises five recognized species: *B. bonitae*, *B. isangdaliri*, *B. ligtas*, *B. mapalanggaon*, and *B. tridactylus*. Of these species, both *B. tridactylus* and true *B. bonitae* are known from a number of sites within the Philippines; however, based on our current understanding of their natural histories, *B. isangdaliri*, *B. mapalanggaon*, and *B. ligtas* have limited ranges (Davis *et al.* 2014; Geheber *et al.* 2016). Due to the secretive and semi-fossorial nature of these skinks, it is likely that additional, undescribed diversity exists within the complex (Davis *et al.* 2014; Geheber *et al.* 2016).

In this paper, we describe a new digitless species of *Brachymeles* in the *B. bonitae* Complex from Luzon Island, discuss its potential distribution, comment on its natural history and ecology, and provide the first illustrations of its diagnostic head scale patterns.

Materials and methods

Field work, sample collection, and specimen preservation. Fieldwork was conducted on Camiguin Norte, Catanduanes, Lubang, Luzon, Marinduque, Masbate, Mindoro, Polillo, Sibuyan, and Tablas islands, all in the Philippines (Fig. 1A), between 1991 and 2012. Specimens were collected during the day, euthanized with MS-222, dissected for tissue samples (liver preserved in 95% ethanol), fixed in 10% formalin, and eventually (< 2 mo) transferred to 70% ethanol. Specimens are deposited in U.S. and Philippine museum collections (see Acknowledgments and Specimens Examined). Museum abbreviations for specimens examined follow those from Sabaj Pérez (2014).

Morphological data. We examined fluid-preserved specimens (Appendix I) for variation in qualitative, meristic (scale counts) and mensural (measurements) characters. Sex was determined by gonadal inspection, and measurements were taken to the nearest 0.1 mm with digital calipers by CDS. X-rays were taken with a company cabinet X-ray on Kodak MIN-R 2000 film exposed at 5 milliamperes and 30 volts for 1 minute 15 seconds.

Meristic and mensural characters were chosen based on Siler *et al.* (2009a, 2010a,b): snout–vent length (SVL), axilla–groin distance (AGD), total length (TotL), midbody width (MBW), midbody height (MBH), tail length (TL), tail width (TW), head length (HL), head width (HW), snout–forearm length (SnFa), eye diameter (ED), eye–nares distance (END), snout length (SNL), fore-limb length (FLL), hind limb length (HLL), midbody scale-row count (MBSR), paravertebral scale-row count (PVSR), axilla–groin scale-row count (AGSR), supralabial count (SL), infralabial count (IFL), supraciliary count (SC), and supraocular count (SO). Additionally, we counted the number of presacral vertebrae (PSV) from X-ray images of specimens. In the description, ranges are followed by mean \pm standard deviation in parentheses.

Distribution model. We estimated the geographic distribution of the new species using ecological niche modeling (ENM) that searches for associations between species' presence and environmental conditions (Soberón & Peterson 2005). We used the SDMtoolbox (v1.1c; Brown 2014) package in ArcMap (v10.2; ESRI 2013) to generate a minimum convex polygon that included the 13 known localities of occurrence for *B. ilocandia* **sp. nov.** and used the polygon to limit the extent of the environmental data input in the model (Barve *et al.* 2011). To characterize the environment, we downloaded 19 bioclimatic variables at 30 arc-seconds resolution (approximately 1 km; Hijmans *et al.* 2005) derived from weather station data from the 1950–2000 period and ran a correlation analysis to remove highly correlated variables, thus limiting collinearity in our dataset. As a result, we retained five variables to run the ecological niche model: annual mean temperature (BIO1), mean diurnal range (BIO2), isothermality (BIO3), annual precipitation (BIO12), and precipitation of driest month (BIO14). All pairs of variables had a correlation coefficient below 0.7, except for two: BIO12 – BIO14 ($|r^2| = 0.8$) and BIO12 – BIO2 ($|r^2| = 0.82$). To generate the niche model and estimate potential distribution, we used the Maxent maximum entropy algorithm (v3.3.3k; Phillips *et al.* 2006, Phillips & Dudík 2008) with the default settings except for the number of

replicates, which we modified from the default (one) to five replicates. Each of these five models was built with a different subset of occurrences split into two groups: one larger subset for model calibration and one smaller subset for model performance evaluation with five-fold cross-validation (Hijmans 2012). The replicates were trained at the geographic extent encompassing all known occurrences for *B. ilocandia* **sp. nov.** and projected at the extent of the entire Philippine archipelago known to possess species of *Brachymeles* to estimate the potential distribution of the species in the region (Brown & Rabor 1967; Siler & Brown 2010; Davis *et al.* 2014). We used the average of Maxent suitability scores of the five replicates to map the potential distribution of the species. To assess the model predictive accuracy of each replicate, we examined omission error, representing the proportion of presences (used for model validation) that are incorrectly predicted absent by the model. Omission error is calculated by converting the continuous Maxent suitability values to presence-absence; all values below the lowest value associated with a presence (from the training dataset; “lowest presence threshold”; Pearson *et al.* 2007) are assigned a value of zero (absent) and all at or above that value are assigned a value of one (present).

Species concept. For this study, we apply the General Lineage Concept of species (de Queiroz 1998, 1999), as an extension of the Evolutionary Species Concept (Simpson 1961; Wiley 1978; Frost & Hillis 1990). We consider phenotypically divergent populations as distinct lineages, especially if such populations are allopatric. In this study we diagnose a new species based on diagnostic character differences in non-overlapping morphological character states.

Research experience in the undergraduate classroom. As part of the Spring 2015 Herpetology Course (BIOL 4083) taught by CDS at the University of Oklahoma, students took part in a semester long, small group writing assignment, with each group assigned a distinct lineage of *Brachymeles* to describe under a structured writing and mentoring program (Siler *et al.* unpublished data). Detailed description of this course project has been made freely available at <http://www.webcitation.org/6hEkRmogM> (Watters & Siler 2016).

Taxonomic account

Brachymeles ilocandia **sp. nov.**

(Figs. 2, 3)

Brachymeles bonitae Duméril & Bibron 1839; Taylor 1917; Brown 1956:5; Brown & Rabor 1967:526; Brown & Alcalá 1970; Brown & Alcalá 1980:20; Davis *et al.* 2014; Geheber *et al.* 2016.

Holotype. PNM 9819 (RMB Field No. 7324, formerly KU 308004), adult female, collected under a decaying log in secondary-growth forest on 10 February 2007, Sitio Kauringan, Barangay Balatubat, Municipality of Calayan, Cagayan Province, Camiguin Norte Island, Philippines (18.902° N, 121.908° E; WGS 84; 303 m elev.), by J. Fernandez.

Paratypes (Paratopotypes). One adult female (KU 307967), collected on 8 February 2006 in Sitio Kauringan, Barangay Balatubat, Municipality of Calayan, Cagayan Province, Camiguin Norte Island, Philippines, by J. Fernandez and RMB. Four adult females, collected under a decaying log in secondary-growth forest on 11 February 2007 (KU 308019, 308020), 12 February 2007 (KU 308027), and 14 February 2007 (KU 308030) in Sitio Kauringan, Barangay Balatubat, Municipality of Calayan, Cagayan Province, Camiguin Norte Island, Philippines, by J. Fernandez and C. Oliveros.

Paratypes. One adult female (CAS 61377), collected in 1920 in Nayon, Ifugao Subprovince, Mountain Province, Luzon Island, Philippines. One adult female (FMNH 259449), collected on 10 March 2001 in Kalinga Province, Luzon Island, Philippines, by RMB and ACD. One adult female (KU 304567), collected on 5 March 2006 in “Limandok,” Barangay Balatubat, Municipality of Calayan, Cagayan Province, Camiguin Norte Island, Philippines (18.92927° N, 121.89881° E; WGS 84; 403 m elev.), by J. Fernandez and RMB.

Diagnosis. Following recent taxonomic revisions of *Brachymeles* (Siler *et al.* 2011; Davis *et al.* 2014; Geheber *et al.* 2016) the new species is assigned to the *B. bonitae* Complex based on the following suite of morphological characters: (1) limbs present, (2) non-pentadactyl, (3) fore-limbs with 0–3 fingers, (4) hind limbs with 0–2 toes, (5) paravertebral scale rows ≥ 91 , (6) presacral vertebrae 47–53, (7) supraoculars four, (8) enlarged, differentiated nuchals present, (9) longitudinal rows of dark spots around the body absent, and (10) auricular opening absent.




FIGURE 2. Illustration of head of *Brachymeles ilocandia* **sp. nov.** (PNM 9819, holotype) in dorsal, lateral, and ventral views. Taxonomically diagnostic head scales are labeled as follows: C, chin shield; F, frontal; FN, frontonasal; FP, frontoparietal; IL, infralabial; IP, interparietal; L, loreal; M, mental; N, nasal; Nu, nuchal; P, parietal; PF, prefrontal; PM, postmental; PO, preocular; PSO, presubocular; R, rostral; SC, supraciliary; SL, supralabial; SN, supranasal; and SO, supraocular. Roman numerals indicate scales in the supraocular series, with Arabic numbers indicating scales in the supraciliary series. Illustrations by MLP and CDS.



FIGURE 3. Photograph of dorsal, lateral, and ventral views of head of the preserved holotype of *Brachymeles ilocandia* **sp. nov.** (PNM 9819). Photographs taken by JLW and MLP.

Brachymeles ilocandia **sp. nov.** can be distinguished from congeners by the following combination of characters: (1) body size small (SVL 65.7–77.6 mm), (2) limbs digitless, (3) limb length short, (4) supralabials six, (5) infralabials five or six, (6) supraciliaries five, (7) supraoculars four, (8) midbody scale rows 22–24, (9) axilla–groin scale rows 80–82, (10) paravertebral scale rows 97–100, (11) mental/first infralabial fusion present or absent, (12) prefrontal contact absent or in point contact, (13) frontoparietal contact present, (14) enlarged chin shields in three pairs, (15) nuchals enlarged, (16) auricular opening absent, (17) presacral vertebrae 50–53, and (18) uniform body color (Tables 1, 2).

TABLE 1. Summary of mensural characters among species of the *Brachymeles bonitae* Complex. Sample size, body length and total length among males and females, and general geographical distribution (PAIC = Pleistocene Aggregate Island Complexes, sensu Brown & Diesmos [2002]) are included for reference (SVL, TotL, FLL, and HLL given as range over mean \pm standard deviation; all body proportions given as percentage over mean \pm standard deviation).

	<i>bonitae</i> (3 m, 1 f)	<i>ilocandia</i> sp. nov. (9 f)	<i>isangdaliri</i> (1 f)	<i>ligtas</i> (3 m, 2 f)	<i>mapalanggaon</i> (3 m, 6 f)	<i>tridactylus</i> (12 m, 9 f)
Range	Luzon & Polillo islands	Luzon & Camiguin Norte islands	Luzon Island	Lubang Island	Masbate Island	West Visayan PAIC
SVL (f)	69.4	65.7–77.6 (73.7 \pm 3.7)	59.5	60.7–69.2 (65.0 \pm 6.0)	61.7–75.8 (67.2 \pm 5.4)	59.9–82.3 (71.4 \pm 6.9)
SVL (m)	69.7–78.4 (72.8 \pm 4.8)	N/A	N/A	69.4–79.6 (74.5 \pm 5.1)	65.1–72.7 (68.4 \pm 3.9)	60.7–77.6 (69.0 \pm 6.0)
TotL (f)	N/A	122.3–146.0 (134.1 \pm 8.0)	106.1	119.4	120.2	133.6
TotL (m)	122.0	N/A	N/A	160.6	112.6–118.6 (115.6 \pm 4.3)	120.9–154.1 (136.0 \pm 9.8)
TL/SVL	73	81–90 (85 \pm 3)	78	97–102 (99 \pm 4)	67–84 (78 \pm 9)	85–112 (95 \pm 10)
FLL	1.0–1.7 (1.3 \pm 0.3)	1.1–1.4 (1.3 \pm 0.1)	1.3	1.2–1.4 (1.3 \pm 0.1)	0.8–1.0 (0.9 \pm 0.1)	1.5–2.5 (2.0 \pm 0.3)
FLL/SVL	1–2 (2 \pm 0)	1–2 (2 \pm 0)	2	2–2 (2 \pm 0)	1–2 (1 \pm 0)	2–4 (3 \pm 0)
HLL	1.5–2.3 (1.9 \pm 0.3)	1.6–1.9 (1.8 \pm 0.1)	2.2	1.6–2.0 (1.8 \pm 0.1)	1.2–1.6 (1.4 \pm 0.1)	2.6–3.6 (3.1 \pm 0.3)
HLL/SVL	2–3 (3 \pm 0)	2–3 (2 \pm 0)	4	2–3 (3 \pm 0)	2–2 (2 \pm 0)	3–6 (5 \pm 1)

Comparisons. *Brachymeles ilocandia* **sp. nov.** can be distinguished from *B. bonitae* by having fewer axilla–groin scale rows (80–82 *versus* 83–90) and fewer paravertebral scale rows (97–100 *versus* 103–110); from *B. ligtas* by having a greater number axilla–groin scale rows (80–82 *versus* 74–76), paravertebral scale rows (97–100 *versus* 91–93), and fewer presacral vertebrae (47 *versus* 50); from *B. isangdaliri* by having fewer supraciliaries (5 *versus* 6), the presence (*versus* absence) of a third chin shield pair, and a longer snout–vent length (65.7–77.6 mm *versus* 59.5); from *B. mapalanggaon* by having a longer fore-limb length (1.1–1.4 mm *versus* 0.8–1.0); from *B. tridactylus* by having a shorter fore-limb length (1.1–1.4 mm *versus* 1.5–2.5); from *B. isangdaliri* and *B. mapalanggaon* by

having a longer total length (122.3–146.0 mm *versus* 106.1 [*B. isangdaliri*] or 120.2 [*B. mapalanggaon*]); from *B. isangdaliri* and *B. tridactylus* by having digitless limbs (*versus* unidactyl [*B. isangdaliri*] or tridactyl [*B. tridactylus*]), a greater number of presacral vertebrae (50–53 *versus* 47 [*B. isangdaliri*, *B. tridactylus*]), greater number of axilla–groin scale rows (80–82 *versus* 73 [*B. isangdaliri*] or 72–79 [*B. tridactylus*]), and a shorter hindlimb length (1.6–1.9 mm *versus* 2.2 [*B. isangdaliri*] or 2.6–3.6 [*B. tridactylus*]).

Brachymeles ilocandia **sp. nov.** can be distinguished from all limbless species of *Brachymeles* by having limbs, and from all pentadactyl species of *Brachymeles* by having digitless limbs.

Description of holotype. Details of the head scalation are shown in Figures 2 and 3. Adult, female, body small, slender, SVL 74.2 mm; head weakly differentiated from neck, nearly as wide as body, HW 5.9% SVL, 86.4% HL; HL 38.4% SnFa; SnFa 17.8% SVL; snout broadly rounded in dorsal and lateral profile, SNL 60.0% HL; ear completely hidden by scales; eyes small, ED 15.9% HL, 42.0% END, pupil subcircular; body slightly depressed, nearly uniform in thickness, MBW 129.1% MBH; scales smooth, glossy, imbricate; longitudinal scale rows at midbody 23; paravertebral scale rows 97; axilla–groin scale rows 80; limbs short, diminutive, bluntly rounded; digits absent; finger and toe lamellae absent; FLL 2.0% AGD, 1.5% SVL; HLL 3.0% AGD, 2.3% SVL; tail not as wide as body, gradually tapered towards end, TW 87.2% MBW, TL 83.3% SVL.

Rostral projecting onto dorsal snout to level in line with midline nasal, roughly equal in width and height, in contact with frontonasal; frontonasal wider than long; nostril ovoid, in posterodorsal corner of single trapezoidal nasal, longer axis directed posterodorsally and anteroventrally; supranasals present; postnasals absent; prefrontals narrowly separated; frontal roughly diamond-shaped, its anterior margin in moderate contact with frontonasal, in contact with first two anterior supraoculars, 4× wider than anterior supraocular; supraoculars four; frontoparietals moderate, moderately separated, each frontoparietal in contact with supraoculars two and three; interparietal large, its length roughly equal to 2× midline length of frontoparietal, longer than wide, diamond-shaped, wider anteriorly, pineal eyespot visible in center; parietals broader than frontoparietals, in broad contact behind interparietal; enlarged nuchals present; loreals two, anterior loreal slightly longer and higher than posterior loreal; preocular one; presubocular one; supraciliaries five, the anteriormost contacting prefrontal and separating posterior loreal from first supraocular, posteriormost extending to midline of fourth supraocular; subocular scale row single, incomplete, in contact with supralabials; lower eyelid with one row of scales; supralabials six, first nearly twice the width of others, third, fourth and fifth subocular; infralabials five (Fig. 2).

Mental wider than long, fused with first infralabials; postmental single, enlarged, its width narrower than width of mental; followed by three pairs of enlarged chin shields, first pair narrowly separated, second pair largest, narrowly separated, third pair smallest, broadly separated by three medial scales (Fig. 2). Scales on limbs smaller than body scales.

Variation. Differences in scalation patterns were observed among the type series. Observed scale row counts variation included: midbody scale rows of 22 (FMNH 259449, KU 308030), 23 (CAS 61377, KU 307967, 308004, 308019, 308027), and 24 (KU 304567, 308020); axilla–groin scale rows of 80 (CAS 61377, FMNH 259449, KU 308004, 308030), 81 (KU 307967, 308019, 308020), and 82 (KU 304567, 308027); paravertebral scale rows of 97 (KU 308004), 98 (FMNH 259449, KU 307967, 308019, 308020, 308030), 99 (CAS 61377), and 100 (KU 304567, 308027); and five (KU 308004, 308027, 308030) or six (CAS 61377, FMNH 259449, KU 304567, 307967, 308019, 308020) infralabials. Additionally, a low level of variation was observed in the presence of a fused mental and first infralabial scale (both sides of head in KU 308004 and 308030, right side of head only in KU 308027). This character was absent in all other specimens examined.

Coloration of holotype in life. Coloration in life is unrecorded; however, because *Brachymeles* specimens do not change significantly during preservation (CDS, RMB *personal observations*), we suspect that the preserved coloration and patterns are much like those in life.

Coloration of holotype in preservative. The dorsal, lateral and ventral regions of the trunk and tail are a solid Prout's Brown (Color 47; Köhler 2012). The head has the same background color, with dense speckling on the dorsal and lateral surfaces, causing the head to appear more Hair Brown (Color 277; Köhler 2012) in color. In the vicinity of each orbit there is a Fuscous (Color 283; Köhler 2012) splotch. The ventral region of the head is the same solid Prout's Brown (Color 47; Köhler 2012) as the trunk.

TABLE 2. Summary of meristic and qualitative diagnostic characters (present, absent) among species of the *Brachymeles bonitae* Complex. The pairs of enlarged scales posterior to the postmental scale are abbreviated as chin shield pairs with reference to the 1st, 2nd, and 3rd pairs (when present). In cases of scale count variation within species, numbers of individuals showing specific counts are given in parentheses. Numbers in parentheses after each species indicate male (m) and female (f) individuals examined.

Diagnostic character	<i>bonitae</i> (3 m, 1 f)	<i>ilocandia</i> sp. nov. (9 f)	<i>isangdaliri</i> (1 f)	<i>ligtas</i> (3 m, 2 f)	<i>mapalanggaon</i> (3 m, 6 f)	<i>tridactylus</i> (12 m, 9 f)
Number of digits (fingers/toes)	0/0 ¹ 2/1 ¹	0/0	1/1	2/0	0/0	3/3
PSV	53	50–53	47	50	51	47
MBSR	21–24	22–24	22	22	22, 23	22–24
AGSR	83–90	80–82	73	74–76	80–84	72–79
PVSR	103–110	97–100	97	91–93	99–102	90–98
SL	6	6	6	6	6	6 (13) 7 (8)
IFL	5	5 (3) 6 (6)	6	6	5 (8) 6 (1)	6 (13) 7 (8)
SC	5	5	6	5	5	5
SO	4	4	4	4	4	4
Prefrontal contact	Absent	Absent or Point contact	Absent	Absent	Absent	Absent
Frontoparietal contact	Absent	Absent	Absent	Present	Point contact or Absent	Absent
1 st chin shield pair contact	Absent	Absent	Absent	Present or Absent	Absent	Present or Absent
3 rd chin shield pair	Present	Present	Absent	Present	Present or Absent	Present
Mental/1 st IFL fusion	Present	Present or Absent	Absent	Absent	Present or Absent	Absent
Enlarged nuchals	Present	Present	Present	Present	Present	Present
Longitudinal rows of dark spots	Absent	Absent	Absent	Absent	Absent	Absent

¹Observed for two individuals.

Distribution, ecology and natural history. *Brachymeles ilocandia* **sp. nov.** is known only from northern Luzon and Camiguin Norte islands (Fig. 1A). The new species likely once occurred in low- to mid-elevation primary forest habitats; however, all recent observations of this species have occurred in secondary growth forest habitats. In contrast to the other members of the *B. bonitae* Complex, this species appears to be quite common in secondary growth forest fragments throughout the northern Philippines. *Brachymeles ilocandia* **sp. nov.** is found in parts of its distribution with *B. bicolor*, *B. kadwa*, and *B. boulengeri*. Other species of *Brachymeles* known to occur in the Luzon PAIC include *B. bicolor*, *B. bonitae*, *B. brevidactylus*, *B. cobos*, *B. elerae*, *B. isangdaliri*, *B. kadwa*, *B. lukbani*, *B. makusog*, *B. minimus*, *B. muntingkamay*, and *B. wrighti* (Davis *et al.* 2014).

We have evaluated this species against the International Union for Conservation of Nature (IUCN) criteria for classification and find that it does not qualify for Critically Endangered, Endangered, Vulnerable, or Near Threatened status. Not only does *B. ilocandia* **sp. nov.** occur on more than one island, but also the species appears common in secondary growth and disturbed habitats throughout its recognized distribution. Therefore, we recommend that this species be classified as Least Concern, LC (IUCN 2015).

Etymology. The specific epithet is chosen in reference to the biogeographically and culturally distinct homeland, “Ilocandia”, of the *Ilokano* people of the northern Philippines, the third largest ethnolinguistic group in the country. Including the Babuyan Island Group north of Luzon Island, Ilocandia stretches from the western coast of northern Luzon, across the Cagayan Valley, to parts of central Luzon and the boundaries of Aurora Province in the east. The region is home to many endemic vertebrates, diverse geographic landscape, unique local cuisine, and rich cultural traditions. Suggested common name: Ilokano Slender Skink.

Discussion

With the description of *Brachymeles ilocandia* **sp. nov.** the total number of species within *Brachymeles* increases to 40. Only 18 species of *Brachymeles* were recognized prior to 2010 (for review, see Siler *et al.* 2010); therefore, described *Brachymeles* diversity has increased by more than 220% within the last decade. Cryptic, as well as phenotypically distinct, evolutionary lineages within *Brachymeles* have been detected recently, largely a result of comprehensive biodiversity surveys across the Philippines, careful examination of vouchered natural history collections, and the use of genetic data (Table 3). Furthermore, it is likely that continued survey efforts throughout the Philippines and thorough investigation of phenotypic and genetic variation will result in the recognition of additional species-level diversity within the genus.

TABLE 3. Uncorrected pairwise sequence divergence (%) for mitochondrial data for focal species of the *Brachymeles bonitae* Complex (*Brachymeles bonitae*, *B. ilocandia* **sp. nov.**, *B. isangdaliri*, *B. ligtas*, *B. mapalanggaon*, and *B. tridactylus*). Percentages on the diagonal represent intraspecific genetic diversity (bolded for emphasis). Data based on Siler *et al.* (2011a) and Davis *et al.* (2014).

	<i>bonitae</i>	<i>ilocandia</i> sp. nov.	<i>isangdaliri</i>	<i>ligtas</i>	<i>mapalanggaon</i>	<i>tridactylus</i>
<i>bonitae</i>	0.2–1.5					
<i>ilocandia</i> sp. nov.	9.5–9.7	0.0				
<i>isangdaliri</i>	9.5–10.0	10.5	0.0			
<i>ligtas</i>	8.8–9.5	6.9–7.0	10.0	0.7		
<i>mapalanggaon</i>	9.0–11.1	10.5–11.2	10.4–11.2	10.3–11.1	2.6	
<i>tridactylus</i>	8.6–9.6	10.5–11.0	9.5–10.3	9.8–10.2	7.8–8.8	0.1–5.0

Brachymeles ilocandia **sp. nov.** is a new species within the *B. bonitae* Complex found in areas of secondary-growth forests on northern Luzon Island and Camiguin Norte Island. Originally thought to be a single widespread species distributed across the central and northern Philippines, *B. bonitae* is now recognized as a species complex containing multiple unique evolutionary lineages (Davis *et al.* 2014; Geheber *et al.* 2016). The new species is the sixth recognized species of the *B. bonitae* Complex, with a unique digitless body form only shared with one other species in the genus (*B. mapalanggaon* from Masbate Island; Davis *et al.* 2014). *Brachymeles ilocandia* **sp. nov.** also possesses an allopatric distribution compared with all other species in the complex (Davis *et al.* 2014; Geheber

et al. 2016), including: true *B. bonitae sensu* Davis *et al.* (2014) from Polillo Island and central Luzon Island, *B. isangdaliri* from southeastern Luzon, *B. mapalanggoan* from Masbate Island, *B. ligtas* from Lubang Island, and *B. tridactylus* from Boracay, Carabao, Inampulugan, Negros, and Panay islands. Lineages of the *B. bonitae* Complex represented in phylogenetic and systematic studies have revealed considerable genetic divergence among populations with available vouchered genetic samples (Siler *et al.* 2011a; Davis *et al.* 2014). In fact, mitochondrial sequence divergences among all recognized members of the complex range from 6.9–11.2%, with an average among lineage uncorrected pairwise mitochondrial sequence divergence of 9.9% (Table 3). This level of divergence is considerable for any species complex, particularly when compared with divergences between previously defined species—viz., *B. cebuensis*, *B. minimus*, *B. lukbani* (Table 3; Siler *et al.* 2011a,b, 2012a; Davis *et al.* 2014).

Interestingly, species within the *B. bonitae* Complex are found on several different PAICs. *Brachymeles ilocandia* **sp. nov.**, *B. bonitae*, and *B. isangdaliri* have been recorded from islands of the Luzon PAIC, *B. mapalanggoan* and *B. tridactylus* from the Western Visayan PAIC, *B. ligtas* from the Mindoro PAIC, and *B. ilocandia* **sp. nov.** from the Babuyan Island Group. Phylogenetic studies focused on the genus have rejected PAIC formation and fragmentation as an explanatory mechanism behind patterns of diversification in this group (Siler *et al.* 2011a); therefore, it is possible that the distribution and diversification of species within the *B. bonitae* Complex occurred as a result of overseas dispersal (Siler *et al.* 2011a). This is further supported by the distribution of *B. ilocandia* **sp. nov.**, which inhabits islands that are part of two separate PAICs and were never connected by land (Fig. 1A).

Furthermore, the results of ecological niche modeling support the possibility that additional envelopes of suitable climate conditions exist across the northeast Luzon PAIC and Babuyan Island Group beyond those verified by vouchered populations of *B. ilocandia* **sp. nov.** observed during biodiversity surveys (Fig. 1B). Specifically, the remainder of the Babuyan Island Group, the entirety of Aurora Province, and eastern Quezon Province (General Nakar Province) on the Luzon PAIC represent regions predicted to possess suitable climatic conditions; however, no records of populations of *B. ilocandia* **sp. nov.** are known from these regions (Fig. 1B). Additional surveys of these regions are necessary to ascertain whether this species truly has a broader range than is currently recognized. These geographic regions are known to possess other species of *Brachymeles*, including microendemic congeners (e.g., *B. elerae* and *B. isangdaliri*; Siler 2010; Siler *et al.* 2011d; Davis *et al.* 2014), and thus represent unique faunal subregions on Luzon Island that warrant further study and conservation efforts (Siler *et al.* 2011d; Davis *et al.* 2014). Moreover, discovery of additional populations of *B. ilocandia* **sp. nov.** in these predicted regions may provide new insights into intraspecific genetic and morphological variation.

Luzon Island is the largest island within the Philippine archipelago, and, as such, contains the greatest number of species of *Brachymeles*. Following the recognition of *B. ilocandia* **sp. nov.**, 13 species are now recognized to occur on Luzon Island: *B. bicolandia*, *B. bonitae*, *B. brevidactylus*, *B. cobos*, *B. elerae*, *B. isangdaliri*, *B. kadwa*, *B. lukbani*, *B. makusog*, *B. minimus*, *B. muntingkamay*, *B. wrighti*, and *B. ilocandia* **sp. nov.** (Davis *et al.* 2014). *Brachymeles ilocandia* **sp. nov.** is co-distributed across parts of its range with the large pentadactyl species *B. bicolor*, *B. kadwa*, and *B. boulengeri* (Siler & Brown 2011; Davis *et al.* 2014). All species of *Brachymeles* are known to be semi-fossorial, with sympatric species often found within the same rotting log or root network (CDS, *personal observation*). However, to date, little is known of the ecology or natural history of species in the genus, and future research on community structure, functional capacities, and species microhabitat requirements within *Brachymeles* should be conducted to improve our understanding of this unique radiation of lizards.

Acknowledgments

We thank the Biodiversity Management Bureau (BMB) of the Philippine Department of Environment and Natural Resources (DENR) for facilitating collecting and export permits necessary for this and related studies; we are particularly grateful to T. M. Lim, C. Custodio, A. Tagtag, and J. L. De Leon for their logistical support of this research. Fieldwork was conducted under the Memorandum of Agreement with the BMB of the Philippines (2009–2014), Gratuitous Permit to Collect No. 221, and KU IACUC Approval (158-01). Financial support for fieldwork was provided by a Panorama Fund grant from The University of Kansas Biodiversity Institute, travel funds from The University of Kansas Ecology and Evolutionary Biology department, a Madison and Lila Self Fellowship from the University of Kansas, a Fulbright Fellowship, a Fulbright-Hayes Fellowship, NSF DEB 0804115 and NSF IOS

1353683 to CDS, and NSF DEB 0743491 and EF-0334952 to R. Brown. For the loans of specimens we thank D. Blackburn, J. Vindum, and A. Leviton (California Academy of Sciences), J. Barnes and A. C. Diesmos (Philippine National Museum), J. Ferner (Cincinnati Museum of Natural History), A. Resetar and H. Voris (Field Museum of Natural History), R. Crombie, K. de Queiroz (National Museum of Natural History, Smithsonian Institution), and D. Cannatella and T. LaDuc (Biodiversity Collections, University of Texas at Austin). For access to the Sam Noble Museum Invertebrate Paleontology Stacking Photography Lab we thank S. Westrop and R. Burkhalter. Critical evaluations of the manuscript and discussion of species concepts were provided by members of the Spring 2015 Herpetology Class at the University of Oklahoma. CDS thanks the CAS Stearns Fellowship and the MCZ Ernst Mayr Fellowship for funding recent visits to examine comparative material. Both CDS and RMB extend a special thanks to A. Alcalá for his continued support of our Philippine biodiversity research program.

References

- Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S.P., Peterson, A.T., Soberón, J. & Villalobos, F. (2011) The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecological Modeling*, 222, 1810–1819.
<http://dx.doi.org/10.1016/j.ecolmodel.2011.02.011>
- Brown, J.L. (2014) SDMtoolbox: a python?based GIS toolkit for landscape genetic, biogeographic and species distribution model analyses. *Methods in Ecology and Evolution*, 5, 694–700.
<http://dx.doi.org/10.1111/2041-210X.12200>
- Brown, R.M. & Diesmos, A.C. (2002) Application of lineage-based species concepts to oceanic island frog populations: the effects of differing taxonomic philosophies on the estimation of Philippine biodiversity. *Silliman Journal*, 42, 133–162.
- Brown, R.M. & Diesmos, A.C. (2009) Philippines, Biology. In: Gillespie, R., Clague, D. (Eds.), *Encyclopedia of Islands*. University of California Press, Berkeley, California, pp. 723–732.
- Brown, R.M. & Guttman, S.I. (2002) Phylogenetic systematics of the *Rana signata* complex of Philippine and Bornean stream frogs: reconsideration of Huxley's modification of Wallace's Line at the Oriental–Australian faunal zone interface. *Biological Journal of the Linnean Society*, 76, 393–461.
<http://dx.doi.org/10.1111/j.1095-8312.2002.tb01704.x>
- Brown, R.M. & Siler, C.D. (2013) Spotted stream frog diversification at the Australasian faunal zone interface, mainland versus island comparisons, and a test of the Philippine dual umbilici hypothesis. *Journal of Biogeography*, 41, 182–195.
<http://dx.doi.org/10.1111/jbi.12192>
- Brown, R.M., Diesmos, A.C. & Alcalá, A.C. (2002) The state of Philippine herpetology and the challenges for the next decade. *Silliman Journal*, 42, 18–87.
- Brown, R.M., Siler, C.D., Diesmos, A.C. & Alcalá, A.C. (2009) The Philippine frogs of the genus *Leptobrachium* (Anura; Megophryidae): phylogeny-based species delimitation, taxonomic revision, and descriptions of three new species. *Herpetological Monograph*, 23, 1–44.
<http://dx.doi.org/10.1655/09-037.1>
- Brown, R.M., Siler, C.D., Oliveros, C.H., Esselstyn, J.A., Diesmos, A.C., Hosner, P.A., Linkem, C.W., Barley, A.J., Oaks, J.R., Sanguila, M.B., Welton, L.J., Blackburn, D.S., Moyle, R.G., Peterson, A.T. & Alcalá, A.C. (2013) Evolutionary processes of diversification in a model island archipelago. *Annual Review of Ecology, Evolution, and Systematics*, 44, 411–435.
<http://dx.doi.org/10.1146/annurev-ecolsys-110411-160323>
- Brown, W.C. (1956) A revision of the genus *Brachymeles* (Scincidae), with descriptions of new species and subspecies. *Breviora*, 54, 1–19.
- Brown, W.C. & Alcalá, A.C. (1970) The zoogeography of the Philippine Islands, a fringing archipelago. *Proceedings of the California Academy of Sciences*, 38, 105–130.
- Brown, W.C. & Alcalá, A.C. (1980) *Philippine Lizards of the Family Scincidae*. Philippine University Press, Dumaguete City, 264 pp.
- Brown, W.C. & Rabor, D.S. (1967) Review of the genus *Brachymeles* (Scincidae), with descriptions of new species and subspecies. *Proceedings of the California Academy of Sciences*, 15, 525–548.
- Davis D.R., Feller, K.D., Brown, R.F. & Siler, C.S. (2014) Evaluating the diversity of Philippine Slender Skinks of the *Brachymeles bonitae* Complex (Reptilia: Squamata: Scincidae): redescription of *B. tridactylus* and descriptions of two new species. *Journal of Herpetology*, 48, 480–494.
<http://dx.doi.org/10.1670/13-173>
- de Queiroz, K. (1998) The general lineage concept of species, species criteria, and the process of speciation: a conceptual unification and terminological recommendations. In: Howard, D.J. & Berlocher, S.H. (Eds.), *Endless Forms: Species and Speciation*. Oxford University Press, New York, pp. 57–75.
- de Queiroz, K. (1999) The general lineage concept of species and the defining properties of the species category. In: Wilson, R.A. (Ed.), *Species: New Interdisciplinary Essays*. Massachusetts Institute of Technology Press, Cambridge,

- Massachusetts, pp. 49–89.
- Diesmos, A.C., Watters, J.L., Huron, N.A., Davis, D.R., Alcalá, A.C., Crombie, R.I., Afuang, L.E., Gee-Das, G., Sison, R.V., Sanguila, M.B., Penrod, M.L., Labonte, M.J., Davey, C.S., Leone, E.A., Diesmos, M.L., Sy, E.Y., Welton, L.J., Brown, R.M. & Siler, C.D. (2015) Amphibians of the Philippines, part I: checklist of the species. *Proceedings of the California Academy of Sciences, Fourth Series*, 62, 457–539.
- Duméril, A.M.C. & Bibron, G. (1839) *Erpétologie Générale ou, Histoire Naturelle Complète des Reptiles*. Librairie Encyclopédique de Roret, Paris, France, 488 pp.
- ESRI (2013) ArcGIS Desktop: Release 10.2. Environmental Systems Research Institute, Redlands, CA.
- Frost, D.R. & Hillis, D.M. (1990) Species in concept and practice: herpetological applications. *Herpetologica*, 46, 87–104.
<http://www.jstor.org/stable/3892607>
- Geheber, A.D., Davis, D.R., Watters, J.W., Penrod, M.L., Feller, K.D., Davey, C.S., Ellsworth, E.D., Flanagan, R.L., Heitz, B.D., Moore, T., Nguyen, M.D.C., Roberts, A., Sutton, J., Sanguila, M.B., Linkem, C.W., Brown, R.M. & Siler, C.D. (2016) Additions to the Philippine slender skinks of the *Brachymeles bonitae* Complex (Reptilia: Squamata: Scincidae: *Brachymeles*) I: a new species from Lubang Island. *Zootaxa*, 4132 (1), 1–14.
<http://dx.doi.org/10.11646/zootaxa.4132.1.1>
- Heyer, W.R. (1972) A new limbless skink (Reptilia: Scincidae) from Thailand with comments on the generic status of the limbless skinks of Southeast Asia. *Feldiana: Zoology*, 58, 109–129.
<http://dx.doi.org/10.5962/bhl.title.3184>
- Hijmans, R.J. (2012) Cross-validation of species distribution models: removing spatial sorting bias and calibration with a null model. *Ecology*, 93, 679–688.
- Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978.
<http://dx.doi.org/10.1002/joc.1276>
- Hikida, T. (1982) A new limbless *Brachymeles* (Sauria: Scincidae) from Mt. Kinabalu, North Borneo. *Copeia*, 1982, 840–844.
<http://dx.doi.org/10.2307/1444095>
- International Union for Conservation of Nature (IUCN) (2015) *IUCN Red List of Threatened Species. Version 2015.4*. Available from: <http://www.iucnredlist.org> (accessed 15 July 2015)
- Köhler, G. (2012) *Color Catalog for Field Biologists*. Herpeton, Offenbach, Germany, 49 pp.
- Linkem, C.W., Hesed, K.M., Diesmos, A.C. & Brown, R.M. (2010) Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). *Molecular Phylogenetics and Evolution*, 56, 572–85.
<http://dx.doi.org/10.1016/j.ympev.2010.03.043>
- Linkem, C.W., Diesmos, A.C. & Brown, R.M. (2011) Molecular systematics of the Philippine forest skinks (Reptilia: Scincidae: *Sphenomorphus*): testing morphological and biogeographic hypotheses of interspecific relationships. *Zoological Journal of the Linnean Society*, 163, 1217–1243.
<http://dx.doi.org/10.1111/j.1096-3642.2011.00747.x>
- Pearson, R.G., Raxworthy, C., Nakamura, M. & Peterson, A.T. (2007) Predicting species' distributions from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. *Journal of Biogeography*, 34, 102–117.
<http://dx.doi.org/10.1111/j.1365-2699.2006.01594.x>
- Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231–259.
<http://dx.doi.org/10.1016/j.ecolmodel.2005.03.026>
- Phillips, S.J. & Dudík, M. (2008) Modeling of species distributions with MaxEnt: new extensions and a comprehensive evaluation. *Ecography*, 31, 161–175.
<http://dx.doi.org/10.1111/j.0906-7590.2008.5203.x>
- Sabaj Pérez, M.H. (Ed.) (2014) *Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an online reference. Version 5.0*. American Society of Ichthyologists and Herpetologists, Washington, D.C. Available from: <http://www.asih.org/> (accessed 1 June 2015)
- Siler, C.D. (2010) Squamata, Scincidae, *Brachymeles elerae* (Taylor, 1917): Rediscovery in Old Balaban, Cordillera Mountain Range, Luzon Island, Philippines, and natural history. *Check List*, 6, 616–618.
<http://www.checklist.org.br>
- Siler, C.D. & Brown, R.M. (2010) Phylogeny-based species delimitation in Philippine slender skinks (Reptilia: Squamata: Scincidae: *Brachymeles*): taxonomic revision of pentadactyl species groups and description of three new species. *Herpetological Monographs*, 24, 1–54.
<http://dx.doi.org/10.1655/HERPMONOGRAPHS-D-10-00003.1>
- Siler, C.D. & Brown, R.M. (2011) Evidence for repeated acquisition and loss of complex body-form characters in an insular clade of Southeast Asian semi-fossorial skinks. *Evolution*, 65, 2641–2663.
<http://dx.doi.org/10.1111/j.1558-5646.2011.01315.x>
- Siler, C.D., Rico, E.L., Duya, M.R. & Brown, R.M. (2009) A new limb-reduced, loam-swimming skink (Squamata: Scincidae: *Brachymeles*) from central Luzon Island, Philippines. *Herpetologica*, 65, 449–459.
<http://dx.doi.org/10.1655/08-076.1>

- Siler, C.D., Diesmos, A.C. & Brown, R.M. (2010a) A new loam-swimming skink, genus *Brachymeles* (Reptilia: Squamata: Scincidae) from Luzon and Catanduanes Islands, Philippines. *Journal of Herpetology*, 44, 49–60.
<http://www.jstor.org/stable/40665006>
- Siler, C.D., Balete, D.S., Diesmos, A.C. & Brown, R.M. (2010b) A new legless loam-swimming lizard (Reptilia: Squamata: Scincidae: genus *Brachymeles*) from the Bicol Peninsula, Luzon Island, Philippines. *Copeia*, 2010, 114–122.
<http://dx.doi.org/10.1643/CH-08-231>
- Siler, C.D., Oaks, J.R., Esselstyn, J.A., Diesmos, A.C. & Brown, R.M. (2010c) Phylogeny and biogeography of Philippine bent-toed geckos (Gekkonidae: *Cyrtodactylus*) contradict a prevailing model of Pleistocene diversification. *Molecular Phylogenetics and Evolution*, 55, 699–710.
<http://dx.doi.org/10.1016/j.ympev.2010.01.027>
- Siler, C.D., Diesmos, A.C., Alcala, A.C. & Brown, R.M. (2011a) Phylogeny of Philippine slender skinks (Scincidae: *Brachymeles*) reveals underestimated species diversity, complex biogeographical relationships, and cryptic patterns of lineage diversification. *Molecular Phylogenetics and Evolution*, 59, 53–65.
<http://dx.doi.org/10.1016/j.ympev.2010.12.019>
- Siler, C.D., Fuiten, A.M., Jones, R.M., Alcala, A.C. & Brown, R.M. (2011b) Phylogeny-based species delimitation in Philippines slender skinks (Reptilia: Squamata: Scincidae) II: taxonomic revision of *Brachymeles samarensis* and description of five new species. *Herpetological Monographs*, 25, 76–112.
<http://dx.doi.org/10.1655/herpmonographs-d-10-00013.1>
- Siler, C.D., Crombie, R.I., Diesmos, A.C. & Brown, R.M. (2011c) Redescription of two poorly known slender skinks, *Brachymeles bicolor* and *Brachymeles pathfinderi* (Reptilia: Squamata: Scincidae), from the Philippines. *Journal of Herpetology*, 45, 355–369.
<http://dx.doi.org/10.1670/10-103.1>
- Siler, C.D., Jones, R.M., Welton, L.J. & Brown, R.M. (2011d) Redescription of tetradactyl, Philippine slender skinks (genus *Brachymeles*). *Herpetologica*, 67, 300–317.
<http://dx.doi.org/10.1655/herpetologica-d-10-00071.1>
- Siler, C.D., Jones, R.M., Diesmos, A.C., Diesmos, M.L. & Brown, R.M. (2012a) Phylogeny-based species delimitation in Philippine slender skinks (Reptilia: Squamata: Scincidae) III: taxonomic revision of the *Brachymeles gracilis* complex and descriptions of three new species. *Herpetological Monographs*, 26, 135–172.
<http://dx.doi.org/10.1655/herpmonographs-d-11-00006.1>
- Siler, C.D., Oaks, J.R., Welton, L.J., Linkem, C.W., Swab, J.C., Diesmos, A.C. & Brown, R.M. (2012b) Did geckos ride the Palawan raft to the Philippines? *Journal of Biogeography*, 39, 1217–1234.
<http://dx.doi.org/10.1111/j.1365-2699.2011.02680.x>
- Siler, C.D., Oaks, J.R., Cobb, K., Ota, H. & Brown, R.M. (2014) Critically endangered island endemic or peripheral population of a widespread species? Conservation genetics of Kikuchi's gecko and the global challenge of protecting peripheral oceanic island endemic vertebrates. *Diversity and Distributions*, 20, 756–772.
<http://dx.doi.org/10.1111/ddi.12169>
- Simpson, G.G. (1961) *Principles of Animal Taxonomy*. Columbia University Press, New York, New York, 247 pp.
- Taylor, E.H. (1917) *Brachymeles*, a genus of Philippine lizards. *The Philippine Journal of Science*, 12, 267–279.
- Watters, J.L. & Siler, C.D. (2016) *Involving Undergraduates in Research – Herpetology, Spring 2015*. Available from: <http://cameronsiler.com/biol4083-docs-2015> (accessed 3 May 2016).
- Welton, L.J., Siler, C.D., Oaks, J.R., Diesmos, A.C. & Brown, R.M. (2013) Multilocus phylogeny and Bayesian estimates of species boundaries reveal hidden evolutionary relationships and cryptic diversity in Southeast Asian monitor lizards. *Molecular Ecology*, 22, 3495–3510.
<http://dx.doi.org/10.1111/mec.12324>
- Welton, L.J., Wood, P.L. Jr., Oaks, J.R., Siler, C.D. & Brown, R.M. (2014) Fossil-calibrated phylogeny and historical biogeography of Southeast Asian water monitors (*Varanus salvator* Complex). *Molecular Phylogenetics and Evolution*, 74, 29–37.
<http://dx.doi.org/10.1016/j.ympev.2014.01.016>
- Wiley, E.O. (1978) The evolutionary species concept reconsidered. *Systematic Zoology*, 21, 17–26.
<http://dx.doi.org/10.2307/2412809>

Author contributions

CDS conceived the idea; CDS, ADG, and JLW carried out assignment instruction and mentoring; ACD, MLD, CHO, RMB, and CDS participated in fieldwork; MLP created scientific illustrations; NAH created the map; MP assisted in creating the map; AA, AA, DC, TH, AM, NP, and LP compiled and analyzed the dataset; AA, AA, DC, TH, AM, NP, and LP led the writing; CDS, DRD, ESF, NAH, ADG, and JLW assisted in finalizing the manuscript for publication; DRD, ESF, NAH, ADG, JLW, MLP, AA, AA, DC, TH, AM, NP, LP, CHO, RMB, and CDS edited drafts of the manuscript.

APPENDIX I. Additional specimens examined.

Specimens examined in addition to those in Geheber *et al.* (2016). Number in parentheses indicates the number of specimens examined. Several sample sizes are greater than those observed in the description due to the examination of sub-adult specimens which were excluded from morphometric analyses.

Brachymeles ilocandia **sp. nov.** (9). CAMIGUIN NORTE ISLAND: CAGAYAN PROVINCE: *Municipality of Calayan*: Holotype (PNM 9819), Paratopotypes (KU 307967, 308019, 308020, 308027, 308030), Paratype (KU 304567); LUZON ISLAND: KALINGA PROVINCE: Paratype (FMNH 259449); LUZON ISLAND: MOUNTAIN PROVINCE: Paratype (CAS 61377).